

BASE CHANGE FOR ELLIPTIC CURVES OVER REAL QUADRATIC FIELDS

LUIS DIEULEFAIT AND NUNO FREITAS

ABSTRACT. Let E be an elliptic curve over a real quadratic field K and F/K a totally real finite Galois extension. We prove that E/F is modular.

1. INTRODUCTION

For F a totally real number field we write $G_F := \text{Gal}(\overline{\mathbb{Q}}/F)$ for its absolute Galois group. For a Hilbert modular form $\mathfrak{f}$ we denote by $\rho_{\mathfrak{f},\lambda}$ its attached λ -adic representation. We say that a continuous Galois representation $\rho : G_F \rightarrow \text{GL}_2(\overline{\mathbb{Q}}_\ell)$ is **modular** if there exists a Hilbert newform $\mathfrak{f}$ and a prime $\lambda \mid \ell$ in its field of coefficients $\mathbb{Q}_{\mathfrak{f}}$ such that we have an isomorphism $\rho \sim \rho_{\mathfrak{f},\lambda}$. In [1] and [2, Section 5], the first named author proved base change for the GL_2 case over $\mathbb{Q}$.

Theorem 1. *Let f be a classical modular form of weight $k \geq 2$ and field of coefficients $\mathbb{Q}_f$. For a prime λ of $\mathbb{Q}_f$ write $\rho_{f,\lambda}$ for the attached λ -adic representation. Let $F/\mathbb{Q}$ be a totally real number field. Then the Galois representation $\rho_{f,\lambda}|_{G_F}$ is (Hilbert) modular in the sense above.*

Let E be an elliptic curve over a real quadratic field K . It is now known that E is modular over K (see [3]). That is, the ℓ -adic representations $\rho_{E,\ell} : G_K \rightarrow \text{GL}_2(\mathbb{Q}_\ell)$ attached to E arise from some Hilbert newform defined over K of parallel weight 2 having rational field of coefficients.

The aim of this note is to prove a base change result for elliptic curves over real quadratic fields. Our main theorem is the following.

Theorem 2. *Let E be an elliptic curve over a real quadratic field K . Let also F/K be a totally real finite Galois extension. Then E/F is modular.*

This result has applications in the context of the Birch and Swinnerton-Dyer conjecture. Indeed, modularity of E after base change guarantees that the L -function $L(E/F, s)$ is holomorphic in $\mathbb{C}$ and, in particular, its order of vanishing at $s = 1$ is a well defined non-negative integer, in agreement with what is predicted by the BSD conjecture. Furthermore, modularity of E/F allows the construction of Stark-Heegner points on E over (not necessarily real) quadratic extensions of F . For details regarding this application we refer the reader to [4] and the references therein.

Acknowledgements. We would like to thank Kęstutis Česnavičius, José María Giral, Victor Rotger and Samir Siksek for their useful comments.

2010 *Mathematics Subject Classification.* Primary 11F80, Secondary 11G05.

Key words and phrases. modularity, elliptic curves, totally real fields, base change.

2. ELLIPTIC CURVES WITH BIG NON-SOLVABLE IMAGE MOD $p = 3, 5$ OR 7

Let F/K be a finite extension of totally real number fields. Let E/K be an elliptic curve. We will say that $\bar{\rho}_{E,p}(G_F)$ is **big** if $\bar{\rho}_{E,p}(G_{F(\zeta_p)})$ is absolutely irreducible, otherwise we say it is **small**. In particular, if $\bar{\rho}_{E,p}(G_F)$ is non-solvable then it is big. We now restate a result from [3].

Theorem 3. *Let $p = 3, 5$ or 7 . Let F/K and E/K be as above. Suppose that $\bar{\rho}_{E,p}(G_F)$ is big. Then E is modular over F .*

The following proposition is well-known.

Proposition 2.1. *Let F/K be a finite Galois extension of totally real fields and E/K an elliptic curve. Let p be any prime and suppose that $\bar{\rho}_{E,p}(G_K)$ is non-solvable. Then $\bar{\rho}_{E,p}(G_F)$ is non-solvable.*

We have the following corollary.

Corollary 2.2. *Let F/K and E/K be as in Proposition 2.1. Let $p = 3, 5$ or 7 . Suppose that $\bar{\rho}_{E,p}(G_K)$ is non-solvable. Then E is modular over F .*

Proof. From the previous proposition we have that $\bar{\rho}_{E,p}(G_F)$ is non-solvable, hence it is big. Thus E/F is modular by Theorem 3. $\square$

3. ELLIPTIC CURVES WITH PROJECTIVE IMAGE S_4 OR A_4 MOD $p = 3, 5$ OR 7

Let E/K be an elliptic curve. We have seen that if $\bar{\rho}_{E,p}$ has big non-solvable image then after a base change to a Galois extension its image is still non-solvable. We now want to understand what can happen when $\bar{\rho}_{E,p}(G_K)$ is big and solvable. We first recall the following well know fact.

Proposition 3.1. *Let E/K be an elliptic curve. Write G for the image of $\bar{\rho}_{E,p}$ in $\mathrm{GL}_2(\mathbb{F}_p)$ and H for its image in $\mathrm{PGL}_2(\mathbb{F}_p)$. Then, there are the following possibilities:*

- (a) G is contained in a Borel subgroup;
- (b) G contains $SL_2(\mathbb{F}_p)$;
- (c) H is cyclic, G is contained in a Cartan subgroup;
- (d) H is dihedral, G is contained in the normalizer of a Cartan subgroup;
- (e) H is isomorphic to A_4 , S_4 or A_5

In cases c), d), e) the prime p does not divide the order of G .

Let $p = 3, 5$ or 7 . Let also G and H be as in the proposition. We divide the cases where $\bar{\rho}_{E,p}(G_K)$ is big and solvable into two types:

- (I) $H \cong S_4$ or A_4 ,
- (II) H is dihedral.

Suppose we are in case (I). Let F/K be a finite Galois extension and set $H_F := \mathbb{P}(\bar{\rho}_{E,p}(G_F))$. We would like that H_F is also isomorphic to A_4 or S_4 since this would mean that $\bar{\rho}_{E,p}(G_F)$ is big and Theorem 3 applies. Since F/K is Galois we have that H_F is normal subgroup of H . Write $I = \{1\}$ for the trivial group and D_4 for the dihedral group in four elements. The normal subgroups of S_4 and A_4 are respectively

- I , D_4 , A_4 and S_4 ,
- I , D_4 and A_4 .

Thus, the cases where Theorem 3 do not apply over F are when the pair of groups (H, H_F) is one of

$$(1) \quad (S_4, D_4), \quad (S_4, I), \quad (A_4, D_4), \quad (A_4, I).$$

Since we are working with totally real fields the complex conjugation has projective image of order 2. Thus the cases with $H_F = I$ cannot happen.

3.1. A Sylow base change. We now deal with the remaining cases from (1). Recall that we want to base change E/K to F where F/K is finite and Galois. Suppose that (H, H_F) is (S_4, D_4) or (A_4, D_4) . Let F_3 be a subfield of F such that the Galois group $\text{Gal}(F/F_3)$ is a 3-Sylow subgroup of $\text{Gal}(F/K)$. In particular, F/F_3 is a solvable extension. We shall shortly prove the following.

Lemma 3.2. *The projective image $H_{F_3} := \mathbb{P}(\bar{\rho}_{E,p}(G_{F_3}))$ is isomorphic to S_4 or A_4 . In particular, $\bar{\rho}_{E,p}(G_{F_3})$ is big.*

From this lemma and Theorem 3 it follows that E/F_3 is modular. Finally, an application of Langlands solvable base change allows to conclude that E/F is modular.

For the proof of Lemma 3.2 we will need the following elementary lemma from group theory.

Lemma 3.3. *Let G be a profinite group. Let $H \subset G$ be subgroup of finite index i . Let N be a normal subgroup of G . Write j for the index of $H/(N \cap H)$ in G/N . Then $j \mid i$.*

Proof. We prove it for the case of finite groups. The required divisibility follows from the following elementary equalities:

$$\begin{aligned} |G| &= |N| \cdot [G : N] \\ |H| &= |N \cap H| \cdot [H : N \cap H] \end{aligned}$$

Dividing the first equality into the second, we conclude that j divides i . □

Proof of Lemma 3.2. Let F_3 be as above and set

$$G := \text{Gal}(\bar{\mathbb{Q}}/K), \quad H := \text{Gal}(\bar{\mathbb{Q}}/F_3), \quad N := \text{Ker}(\mathbb{P}\bar{\rho}_{E,p}).$$

Let L/K be the Galois extension fixed by N . Observe that $L/L \cap F_3$ is Galois and

$$G/N \cong \text{Gal}(L/K), \quad H/(H \cap N) \cong \text{Gal}(L/L \cap F_3).$$

From Lemma 3.3 we see that

$$[\text{Gal}(L/K) : \text{Gal}(L/L \cap F_3)] = j \mid i = [G : H]$$

and we also have

$$|\text{Gal}(L/K)| = j |\text{Gal}(L/L \cap F_3)|.$$

Note that $\text{Gal}(L/L \cap F_3) \cong H_{F_3}$. From the way we choose F_3 it is clear that $3 \nmid i$, hence $3 \nmid j$. By hypothesis $G/N \cong S_4$ or A_4 , hence 3 divides $|\text{Gal}(L/K)|$ and $|H_{F_3}|$. Finally, the conditions $3 \mid |H_{F_3}|$ and $D_4 \subset H_{F_3}$ together imply that H_{F_3} is isomorphic to S_4 or A_4 . □

We summarize this section into the following corollary.

Corollary 3.4. *Let F/K be a finite Galois extension of totally real fields. Let E/K be an elliptic curve. Suppose that for $p = 3, 5$ or 7 we have that $\bar{\rho}_{E,p}(G_K)$ is big and solvable. Suppose further that $\mathbb{P}(\bar{\rho}_{E,p}(G_K)) \cong S_4$ or A_4 . Then E/F is modular.*

Everything we have done so far works for any Galois extension F/K . Moreover, it is clear that the remaining cases are those when $\bar{\rho}_{E,p}(G_K)$ is small or projectively dihedral simultaneously for $p = 3, 5, 7$. The restriction in the statement of Theorem 2 to quadratic fields arises precisely from dealing with them, which is the content of the next section.

4. ELLIPTIC CURVES HAVING SMALL OR PROJECTIVE DIHEDRAL IMAGE AT $p = 3, 5$ AND 7

Let K be a real quadratic field. From Theorem 3 an elliptic curve E/K is modular over K except possibly if $\bar{\rho}_{E,p}(G_K)$ is small simultaneously for $p = 3, 5, 7$. Suppose $K \neq \mathbb{Q}(\sqrt{5})$. In [3] it is shown that such an elliptic curve gives rise to a K -point on one of the following modular curves:

$$X(b5, b7), \quad X(b3, s5), \quad X(s3, s5), \\ X(b3, b5, d7), \quad X(s3, b5, d7), \quad X(b3, b5, e7), \quad X(s3, b5, e7),$$

where b, s and ns respectively stand for ‘Borel’, ‘normalizer of split Cartan’ and ‘normalizer of non-split Cartan’. The notation d7 and e7 is explained in [3, section 10], here we remark only that they indicate mod 7 level structures that are respectively finer than ‘normalizer of split Cartan’ and ‘normalizer of non-split Cartan’. Denote by $\mathcal{E}_K$ the set of elliptic curves (up to quadratic twist) corresponding to K -points in the previous modular curves.

In [3] it is also shown that an elliptic curve $E/\mathbb{Q}(\sqrt{5})$ with simultaneously small image for $p = 3, 5, 7$ gives rise to a $\mathbb{Q}(\sqrt{5})$ -point in one of the following modular curves

$$X(d7), \quad X(e7), \quad X(b3, b7), \quad X(s3, b7).$$

Denote by $\mathcal{E}_{\mathbb{Q}(\sqrt{5})}$ the set of elliptic curves (up to quadratic twist) corresponding to $\mathbb{Q}(\sqrt{5})$ -points in these four modular curves.

Furthermore, it also follows from [3] that, for any real quadratic field K , we have

- (i) $\mathcal{E}_K$ contains all elliptic curves (up to quadratic twist) with small or projective Dihedral image simultaneously at $p = 3, 5, 7$;
- (ii) $\mathcal{E}_K$ is finite;
- (iii) Let $E \in \mathcal{E}_K$. Then, either E is a $\mathbb{Q}$ -curve or E has complex multiplication or $\bar{\rho}_{E,7}(G_K)$ contains $\mathrm{SL}_2(\mathbb{F}_7)$.

We can now easily prove the following.

Corollary 4.1. *Let K be a real quadratic field. Let $E \in \mathcal{E}_K$. Let F/K be a finite Galois extension. Then E/F is modular.*

Proof. From (iii) above we know that either (a) E/K is a $\mathbb{Q}$ -curve or has complex multiplication or (b) $\rho_7(G_K)$ is non-solvable. In case (a) base change is already known. In case (b), it follows from Corollary 2.2 that E/F is modular. $\square$

5. PROOF OF THE MAIN THEOREM

Let K be a real quadratic field and E/K an elliptic curve. Write $\bar{\rho}_p = \bar{\rho}_{E,p}$. The curve E/K must satisfy at least one of the following three cases

- (1) $\bar{\rho}_p(G_K)$ is big and non-solvable for some $p \in \{3, 5, 7\}$,

- (2) $\bar{\rho}_p(G_K)$ is big, solvable and satisfy $\mathbb{P}(\bar{\rho}_p(G_K)) \cong S_4, A_4$ for some $p \in \{3, 5, 7\}$,
- (3) E/K belongs to the set $\mathcal{E}_K$.

Let F/K be a totally real finite Galois extension. In each case, modularity of E/F now follows directly from one of the previous sections:

Case (1): this is Corollary 2.2.

Case (2): this is Corollary 3.4.

Case (3): this is Corollary 4.1. □

REFERENCES

- [1] L. Dieulefait, *Automorphy of $\text{Sym}^5(GL(2))$ and base change (with Appendix A by R. Guralnick and Appendix B by L. Dieulefait and T. Gee)*, (submitted)
- [2] L. Dieulefait, *Langlands Base Change for $GL(2)$* , *Annals of Math.* **176** (2012), 1015–1038.
- [3] N. Freitas, B. V. Le Hung and S. Siksek, *Elliptic curves over real quadratic fields are modular*, [arXiv:1310.7088](https://arxiv.org/abs/1310.7088), 13 November 2013.
- [4] X. Guitart, V. Rotger and Y. Zhao, *Almost totally complex points on elliptic curves*, *Trans. AMS*, to appear.

MATHEMATISCHES INSTITUT, UNIVERSITÄT BAYREUTH, 95440 BAYREUTH, GERMANY
E-mail address: `nunobfreitas@gmail.com`

E-mail address: `ldieulefait@ub.edu`